

Chronic Liver Disease And Cirrhosis Deaths

Chronic Liver Disease And Cirrhosis Deaths

- Most recent data (2012): 9.7 deaths per 100,000, including 209 deaths.
- Trend over time: 2000-12: flat
- Description: Includes deaths from alcoholic liver disease, chronic hepatitis, and cirrhosis of liver

Chronic liver disease and cirrhosis deaths, King County, 2008-2012 average

	Rate	Lower CI	Upper CI	Count per Year
King County (all ages)	8.6	8.1	9.2	180
AGE				
<18	0.0	0.0	0.2	0
18-24	0.0	0.0	0.4	0
25-44	3.3	2.6	4.0	20
45-64	21.9	20.1	23.8	113
65+	22.4	19.7	25.5	47
GENDER				
Male	11.3	10.4	12.3	114
Female	6.2	5.5	6.9	67
RACE/ETHNICITY				
AIAN	38.9	27.2	57.5	7
Asian	2.5	1.7	3.5	6
Black	7.8	5.5	10.8	9
Hispanic	7.6	5.2	11.5	8
Multiple	5.1 §	2.4 §	11.7 §	2
NHPI	*	*	*	*
White	9.4	8.7	10.1	154
NEIGHBORHOOD POVERTY				
High	15.2	13.1	17.6	38
Medium	8.6	7.9	9.4	113
Low	5.4	4.5	6.4	28
REGION				
East	5.6	4.7	6.6	31
North	8.3	6.3	10.8	12
Seattle	9.1	8.1	10.3	58
South	10.5	9.5	11.6	78

Comparisons:

- Higher than King County rate (yellow)
- Lower than King County rate (green)

Notes:

Source: WA State DOH, Center for Health Statistics, Death Certificates.

Prepared by Public Health - Seattle & King County, APDE, 01/2015.

*Rate = cases per 100,000 population, age-adjusted to the 2000 US population.


CI is 95% Confidence Interval.

* Too few cases to protect confidentiality and/or report reliable rates.

§ Too few cases to meet precision standard, interpret with caution.

Persons of Hispanic ethnicity can be of any race and are included in the racial categories.

Chronic liver disease and cirrhosis deaths King County, 2008-2012 average



Source: WA State DOH, Center for Health Statistics, Death Certificates.

Prepared by Public Health - Seattle & King County, APDE, 01/2015.

[^]Rate = cases per 100,000 population, age-adjusted to the 2000 US population.

[-----] Confidence interval shows range that includes true value 95% of the time.

^{*} Too few cases to protect confidentiality and/or report reliable rates.

[§] Too few cases to meet precision standard, interpret with caution.

Persons of Hispanic ethnicity can be of any race and are included in the racial categories.

Poverty = Neighborhood poverty levels defined by median household income.

Chronic liver disease and cirrhosis deaths by comparison areas

There is no Healthy People 2020 Objective for chronic liver disease and cirrhosis deaths.

	Latest Available Year	Rate	Lower CI	Upper CI
United States	2010	9.4	N/A	N/A
WA State	2012	11.4	10.6	12.2
King County	2012	9.7	8.4	11.1

Notes:

Source: WA State DOH, Center for Health Statistics, Death Certificates.


Prepared by Public Health - Seattle & King County, APDE, 01/2015.

*Rate = cases per 100,000 population, age-adjusted to the 2000 US population.

N/A: Not available.

CI is 95% Confidence Interval.

Chronic liver disease and cirrhosis deaths by comparison areas



Source: WA State DOH, Center for Health Statistics, Death Certificates.
Prepared by Public Health - Seattle & King County, APDE, 01/2015.
[^]Rate = cases per 100,000 population, age-adjusted to the 2000 US population.
|-----| Confidence interval shows range that includes true value 95% of the time.

Chronic liver disease and cirrhosis deaths by health reporting areas, King County, 2008-2012 average

	Rate	Lower CI	Upper CI	Count per Year
King County (all ages)	8.6	8.1	9.2	180
Auburn	16.2	12.1	21.4	11
..Auburn-North	13.2	8.6	19.6	5
..Auburn-South	20.6	13.6	30.6	6
Bear Creek/Carnation/Duvall	5.6	3.2	9.9	4
Bellevue	5.7	4.1	7.8	9
..Bellevue-Central	7.8	4.1	13.6	3
..Bellevue-NE	5.8	3.0	10.5	2
..Bellevue-South	4.6 §	2.2 §	9.8 §	2
..Bellevue-West	4.5 §	1.9 §	9.9 §	2
Black Diamond/Enumclaw/SE County	7.2	4.3	11.7	4
Bothell/Woodinville	12.2	7.4	19.4	4
Burien	16.6	12.1	22.4	9
Covington/Maple Valley	6.1	3.5	10.4	4
Des Moines/Normandy Park	13.2	8.7	19.5	6
East Federal Way	7.7	4.1	13.8	3
Fairwood	6.2 §	2.2 §	15.0 §	1
Federal Way	10.8	7.9	14.5	10
..Fed Way-Central/Military Rd	11.1	7.5	16.1	6
..Fed Way-Dash Point/Woodmont	10.4	6.0	17.4	4
Issaquah	5.9 §	2.8 §	11.9 §	2
Kenmore/LFP	5.4	2.8	10.1	2
Kent	8.5	6.2	11.4	10
..Kent-East	9.6	5.5	16.3	3
..Kent-SE	9.2	5.8	14.1	5
..Kent-West	5.1 §	2.0 §	12.6 §	1
Kirkland	7.4	4.9	10.7	6
..Kirkland	6.9	4.0	11.5	3
..Kirkland North	8.2	4.3	14.7	3
Mercer Isle/Pt Cities	3.6 §	1.0 §	10.6 §	1
Newcastle/Four Creeks	9.3	4.9	16.8	3
North Highline	11.0 §	5.2 §	22.0 §	2
Redmond	4.4 §	2.2 §	8.4 §	2
Renton	9.5	7.1	12.6	10
..Renton-East	3.0 §	1.0 §	8.5 §	1
..Renton-North	8.3	4.2	15.0	2
..Renton-South	14.1	9.8	19.8	7
Sammamish	*	*	*	*
SeaTac/Tukwila	15.7	11.0	22.2	7
Seattle	9.1	8.1	10.3	58
..Ballard	9.9	6.4	15.1	5
..Beacon/Gtown/S.Park	7.3	4.2	12.2	3
..Capitol Hill/E.lake	7.8	4.3	14.4	3
..Central Seattle	5.7	3.0	10.4	3
..Delridge	9.9	5.5	17.5	3
..Downtown	27.8	21.4	42.4	13
..Fremont/Greenlake	5.2	2.6	10.5	2
..NE Seattle	5.7	3.3	9.5	3
..North Seattle	6.3	3.5	11.2	3
..NW Seattle	10.2	6.5	16.0	5
..QA/Magnolia	6.0	3.5	10.4	4
..SE Seattle	8.5	5.0	13.8	4
..West Seattle	10.3	7.1	15.1	7
Shoreline	8.1	5.3	12.2	6
Snoqualmie/North Bend/Skykomish	7.6	4.0	14.0	3
Vashon Island	6.6 §	2.3 §	22.8 §	1

Comparisons:

 Higher than King County rate (yellow)

 Lower than King County rate (green)

Notes:

Source: WAState DOH, Center for Health Statistics, Death Certificates.

Prepared by Public Health - Seattle & King County, APDE, 01/2015.

*Rate = cases per 100,000 population, age-adjusted to the 2000 US population.


CI is 95% Confidence Interval.

* Too few cases to protect confidentiality and/or report reliable rates.

§ Too few cases to meet precision standard, interpret with caution.


Persons of Hispanic ethnicity can be of any race and are included in the racial categories.

Chronic liver disease and cirrhosis deaths by health reporting areas King County, 2008-2012 average



Note: HRA labels on the map match the chart on the right, listed in alphabetical order
 Source: WA State DOH, Center for Health Statistics, Death Certificates.
 Prepared by Public Health - Seattle & King County, APDE, 01/2015.
 ^Rate = cases per 100,000 population, age-adjusted to the 2000 US population.

**Chronic liver disease and cirrhosis deaths
King County, 2000-2012, three-year rolling averages**



Source: WA State DOH, Center for Health Statistics, Death Certificates.
 Prepared by Public Health - Seattle & King County, APDE, 01/2015.
 ^Rate = cases per 100,000 population, age-adjusted to the 2000 US population.